

Prostate Biopsy Preparation

Patient should be advised to contact the office **immediately** if he/she is taking **Coumadin**.

Discontinue 10 days prior to treatment: *Ticlid*

Discontinue 7 days prior to treatment:

Aspirin (All brands, including baby Aspirin)

All aspirin containing medications

Buffered aspirin (All Brands)

Alka-Seltzer	Equagesic	Panodynes
Argesic-SA	Fiorinal (Butalbital Compound)	Pepto-Bismol
Artha-G	Gaysal-S	Percodan (Oxycodone and Aspirin)
Arthropan	Gemisyn	Persistin
Asproject	Lanorinal	Plavix
Axotal	Lobac	Relafen (Nabumetone)
Buff A	Lortab-ASA	Robaxisal
Buffets II	Magan	Rowasa
Butazolidin (Phenylbutazone)	Magsal	Sal-Fayne
Coricidin	Marnal	Salatin
CP-2	Micrainin	Saleto
Darvon (Propoxyphene Compound)	Midol	Salflex
Dasin	Mobidin	Sine-Off (Aspirin Formula)
Daypro (Oxaprozin)	Mobigesic	Soma-Compound
Disalcid	Momentum	Supac
Doans Pills	Monogesic	Synalgos
Dolcin	Neocyclate	Talwin Compound
Double-A	Norgesic	Trental
Duoprin	Oxalid	Trigesic
Duradyne	Pabalate	Trilisate
Dynosal	PAC	Uracel-S
Emagrin	Panasal 5/500	Vanquish

Discontinue 3 Days Prior to Treatment:

Naproxen (Aleve, Anaprox, Naprosyn, Naprelen)

Actron (Ketoprofen)

Lodine (Etodolac)

Oruvail (Ketoprofen)

Feldene (Piroxicam)

Orudis (Ketoprofen)

Voltaren (Diclofenac Sodium)

Discontinue 2 Days Prior to Treatment:

Ibuprofen (Motrin, Advil, Nuprin, Rufen)

Ansaid (Flurbiprofen)

Dolobid (Diflunisal)

Persantine (Dipyridole)

Anturane (Sulfinpyrazone)

Indocin (Indomethacin)

Ponstel (Mefenamic Acid)

Arthrotec (Diclofenac)

Meclomen
(Meclofenamate)

Tolectin (Tolmetin)

Cataflam (Diclofenac)

Medaprin

Toradol (Ketorolac)

Clinoril (Sulindac)

Nalfon (Fenoprofen)